



ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Nuclear Physics A 758 (2005) 276c–279c

NUCLEAR
PHYSICS **A**

The Search for Supernova Signatures in an Ice Core

A. L. Cole^{a*}, R. N. Boyd^b, M. E. Davis^c, L. G. Thompson^c, A. M. Davis^{d e}, R. S. Lewis^e, E. Zinner^f

^aNational Superconducting Cyclotron Laboratory, Michigan State University, East Lansing, MI 48824

^bPhysics and Astronomy Departments, The Ohio State University, Columbus, OH 43210

^cByrd Polar Research Center and Department of Geological Sciences, The Ohio State University, Columbus, OH 43210

^dDepartment of the Geophysical Sciences, University of Chicago, Chicago, IL 60637

^eEnrico Fermi Institute, University of Chicago, Chicago, IL 60637

^fLaboratory for Space Sciences and Physics Department, Washington University, St Louis, MO 63130

It has been suggested that ice cores may preserve detectable enhancements of some terrestrially rare radioisotopes, ^{10}Be , ^{26}Al , ^{36}Cl , resulting from a near-Earth core-collapse supernova(SN) [1]. Both ^{10}Be and ^{36}Cl are also produced by atmospheric cosmic ray spallation and hence are influenced by processes that modulate the Earth's cosmic ray flux. Previous studies [2], [3] have suggested that enhancements occurred in the ^{10}Be and ^{36}Cl fluxes at ~ 35 ky and at ~ 60 ky for ^{10}Be . Thus we have searched for potential SN condensates with ^{26}Al amongst grains filtered from the 308.6m Guliya ice core recovered from the Qinghai-Tibetan plateau in China [3].

We searched for potential core-collapse SN condensate grains corundum (Al_2O_3), hibonite ($\text{CaAl}_{12}\text{O}_{19}$) and spinel (MgAl_2O_4) (see [4]) in Guliya grain samples from the following time periods: ~ 2 -10 ky, ~ 25 -27 ky, ~ 34 -36 ky, ~ 53 -57 ky, ~ 59 -62 ky and ~ 68 -72 ky. These minerals are rare among terrestrial rocks and fine-grained atmospheric dust of terrestrial origin. Furthermore, they are insoluble in the acids employed in the sample preparation process and therefore separable from other minerals, such as silicates, that have high terrestrial abundances. Candidate SN condensate grains were identified among their terrestrial diluents employing a procedure developed at the University of Chicago for detecting presolar grains in meteoritic samples [5]. A set of 37 grains from the ~ 34 -36 ky, ~ 53 -57 ky and ~ 59 -62 ky samples were analyzed with the NanoSIMS at Washington University to measure their oxygen isotopic ratios. The preliminary results indicate that

*The authors acknowledge the assistance of S. Amari in sample preparation. This work is supported in part by NSF grants PHY-0110253, PHY-0099476 and by NASA grants NAG5-12297, NAG5-11903 and NAG5-11545.

the analyzed grains, representing < 15% of those identified, do not possess the extreme O isotopic ratios expected to characterize a SN source [6], [7].

1. Sample Preparation

The six Guliya grain samples were size sorted into a coarse fraction, $> 2\mu\text{m}$ diameter, and a fine fraction, $\leq 2\mu\text{m}$. Approximately half of the fine fraction, representing $\sim 10^8$ grains per sample, was employed in the analysis, with the remaining grains being saved to preserve the archive and the prospect of future analysis. Following the size sorting, targeted chemical dissolutions of unwanted minerals were used to reduce the number of grains to a quantity, on the order of 10^3 , that could be mounted and examined by Scanning Electron Microscope (SEM) Energy Dispersive Spectroscopy (EDS) for candidate grains [5]. Finally, the grains were transferred from suspension onto high purity gold foils with the aim of mounting no more than 2,000 to 5,000 grains. We focused our analysis on grains of diameter $\leq 2\mu\text{m}$, for two reasons. First, the majority of grains in the fine fraction size distribution would have diameters less than $< 10\mu\text{m}$, which is the size capable of surviving descent through the Earth's atmosphere with a speed of 20 km/s as described in [8]. Second, smaller grains experience less atmospheric heating and oxygen isotopic exchange during entry than larger grains.

2. Elemental Mapping

Each of the sample mounts was covered by a grid composed of approximately 300 $100\mu\text{m} \times 125\mu\text{m}$ frames. Elemental X-ray maps were made of each frame of the samples to identify the refractory grains spinel, hibonite and corundum. The number of frames with corundum and other refractory grains identified among the six samples were: ~ 2 -10 ky, 102 frames with corundum; ~ 25 -27 ky, 1 spinel grain and 211 frames with corundum; ~ 34 -36 ky, 69 frames with corundum; ~ 53 -57 ky, 134 frames with corundum; ~ 59 -62 ky, 3 spinel grains, 1 hibonite grain and 70 frames with corundum; ~ 68 -72 ky, 2 spinel grains and 105 frames with corundum. We present the number of frames that possessed Al_2O_3 grains, as some of these frames contained multiple corundum grains and/or aggregates of on the order 10^2 corundum grains that were difficult to enumerate.

3. NanoSIMS

The oxygen isotopic compositions of 37 grains from the ~ 34 -36 ky, ~ 53 -57 ky and ~ 59 -62ky samples were measured with the NanoSIMS at Washington University; a more detailed description of NanoSIMS analysis can be found in [9]. Figure 1 displays the measured $\delta^{18}\text{O}$, $\delta^{17}\text{O}$ ratios for each sample, the associated 1σ errors, the terrestrial fraction (TF) line, and the calcium aluminum inclusion (CAI) mixing line. Before the analysis, terrestrial Al_2O_3 grains were placed on each of the three gold mounts to serve as standards. Following the method of [9], we assume that the oxygen isotope ratios of these terrestrial Al_2O_3 grains are equivalent to standard mean ocean water (SMOW) values.

As shown in Figure 1, five of the eleven Al_2O_3 grains from the ~ 34 -36 ky sample are within 1σ of the CAI mixing line with $\delta^{18}\text{O} \sim -40$ per mil and $\delta^{17}\text{O} \sim -60$ per mil. Two of the eleven grains reside below the CAI mixing line by more than 1σ . All ten ~ 53 -57

ky grains are roughly within 2σ of the TF line and near $\delta^{18}\text{O}\sim 0$ per mil and $\delta^{17}\text{O}\sim 0$ per mil. Finally, the isotopic ratios of 13 Al_2O_3 and three MgAl_2O_4 grains from the ~ 59 –62 ky sample were measured. Two of the spinel grains reside within 1σ of the CAI mixing line at $\delta^{18}\text{O}\sim -40$ per mil and $\delta^{17}\text{O}\sim -40$ per mil. These two spinel grains are located in the same $100\mu\text{m}\times 125\mu\text{m}$ frame and may represent two pieces of a single grain that was fractured during the mounting procedure.

4. Discussion

Previous studies [2], [3] have indicated that enhancements occurred in the ^{10}Be and ^{36}Cl fluxes at ~ 35 ky and at ~ 60 ky for ^{10}Be . None of the analyzed corundum and spinel grains from the time periods ~ 34 –36 ky, ~ 53 –57 ky, ~ 59 –62 ky possessed the anomalous, non-CAI values of $\delta^{18}\text{O}$, $\delta^{17}\text{O}$ indicative of ^{16}O enhancements expected from a core-collapse SN source [6], [7]. However, the 37 grain set represents $< 15\%$ of the total number identified from the elemental maps. The NanoSIMS results indicate that seven corundum grains from the ~ 34 –36 ky sample and two spinel grains from the ~ 59 –62 ky sample possess $\delta^{18}\text{O}$, $\delta^{17}\text{O}$ values that suggest ^{16}O enhancements consistent with CAIs which are typically found in carbonaceous chondrites [10]. These grains are likely high temperature condensates from the solar nebulae. As the grains are in the fine fraction $< 2\mu\text{m}$, they are unlikely to have experienced significant size alteration during the atmospheric entry. The detected CAI grains represent $\sim 1/3$ of the 37 grains analyzed and the number detected differs between the three samples with the majority residing in the ~ 34 –36 ky epoch. Additional analysis of grains from the above samples, ~ 34 –36 ky, ~ 53 –57 ky, ~ 59 –62 ky, and of the remaining three samples, ~ 2 –10 ky, ~ 25 –27 ky, ~ 68 –72 ky, is needed to determine if the variation is the result of the small number of grains sampled or an indication of a time variation in the number of CAI grains accreted by the Earth.

REFERENCES

1. J. Ellis, B.D. Fields and D.N. Schramm, *Astrophys. J.* 470 (1996) 1227.
2. G.M. Raisbeck, F. Yiou, D. Bourles, C. Lorius, J. Jouzel and N. I. Barkov, *Nature* 326 (1987) 273.
3. L.G. Thompson, T. Yao, M.E. Davis, K.A. Henderson, E. Mosley-Thompson, P.-N. Lin, J. Beer, H.-A. Synal, J. Cole-Dai and J.F. Bolzan, *Science* 276 (1997) 1821.
4. D.S. Ebel and L. Grossman, *Geochim. Cosmochim. Acta* 65 (2001) 469.
5. S. Amari, R.S. Lewis and E. Anders, *Geochim. Cosmochim. Acta* 58 (1994) 459.
6. S. Amari and E. Zinner, *Nucl. Phys. A* 621 (1997) 99c.
7. T. Rauscher, A. Heger, R.D. Hoffman and S.E. Woosley, *Astrophys. J.* 576 (2002) 323.
8. S.G. Love and D.E. Brownlee, *Icarus* 89 (1991) 26.
9. E. Zinner, S. Amari, R. Guinness, A. Nguyen, F. J. Stadermann, R.M. Walker and R.S. Lewis, *Geochim. Cosmochim. Acta* 67 (2003) 5083.
10. R.N. Clayton, *Annu. Rev. Earth Planet Sci.* 21 (1993) 115.




Figure 1. The oxygen isotope ratios and 1σ errors of refractory oxide grains from three Guliya ice core grain samples measured with a NanoSIMS.